



THE IMPORTANCE OF COMPREHENSIVE PATIENT EXAMINATION IN DENTAL IMPLANTATION

Khotamov Olloyor Adkham ugli

*Resident Doctor — CAMU (Central Asian Medical
University)*

ARTICLE INFO:

Received:06.10.2026

Revised: 07.10.2026

Accepted:08.10.2026

KEY WORDS:

*dental implantation,
comprehensive
examination,
implantology, patient
assessment, diagnostics,
cone-beam computed
tomography, bone
quality,
osseointegration,
treatment planning,
implant complications.*

ABSTRACT

Dental implantation is one of the most effective and widely used methods for restoring missing teeth and improving oral function and aesthetics. The long-term success of implant treatment largely depends on the accuracy of diagnosis and the quality of preoperative examination. A comprehensive patient assessment makes it possible to identify systemic diseases, local oral conditions, anatomical limitations, bone quality, periodontal status, and other risk factors that may affect osseointegration and implant survival. Modern diagnostic methods, including clinical examination, radiographic assessment, cone-beam computed tomography, laboratory investigations, and evaluation of the patient's medical history, allow the clinician to create an individualized treatment plan.

Dental implants have become an important part of modern restorative dentistry because they provide a functional and aesthetic solution for patients with partial or complete tooth loss. However, successful implant therapy cannot be achieved only through correct surgical technique. The condition of the patient's general health, oral cavity, bone tissue, periodontal structures, and anatomical features must also be carefully evaluated before surgery. According to contemporary implantology principles, detailed preoperative examination is one of the main factors influencing implant survival and long-term stability [1]. A comprehensive examination begins with obtaining a detailed medical and dental history. Diseases such as uncontrolled diabetes mellitus, cardiovascular disorders, osteoporosis, immune system disorders, and certain systemic conditions may negatively influence wound healing and osseointegration. In addition,



smoking, poor oral hygiene, alcohol consumption, and long-term medication use can increase the risk of implant failure and peri-implant complications [2]. For this reason, the clinician should assess not only the local condition of the oral cavity but also the general health status of the patient.

Clinical examination includes evaluation of the teeth, gingiva, periodontal tissues, occlusion, interarch relationships, mucosal condition, and the amount of available keratinized tissue. The presence of untreated caries, periodontal disease, infection, or poor oral hygiene may increase the risk of postoperative complications. Periodontal health is especially important because patients with a history of periodontitis may have a higher risk of developing peri-implantitis after implant placement [3]. Therefore, all active infections and periodontal problems should be treated before surgical intervention.

Radiological examination is another essential part of pre-implant diagnosis. Conventional panoramic radiography can provide general information about the jaws; however, cone-beam computed tomography is considered particularly valuable for three-dimensional assessment of the alveolar bone and nearby anatomical structures. CBCT helps determine bone height, width, density, and the exact position of structures such as the maxillary sinus, nasal cavity, and inferior alveolar nerve [4]. This information allows the clinician to select the optimal implant length, diameter, position, and angulation while reducing the risk of damaging important anatomical structures.

The effectiveness of dental implantation is closely connected with correct patient selection and accurate treatment planning. Comprehensive examination makes it possible to identify possible contraindications before surgery and to modify the treatment plan according to individual risk factors. For example, patients with insufficient bone volume may require bone augmentation, sinus lifting, or other regenerative procedures before implant placement. Similarly, patients with poorly controlled systemic diseases may need medical consultation and stabilization of their condition before surgical treatment.

Bone quality and quantity play a central role in achieving primary implant stability. When the clinician has reliable information about the structure of the alveolar ridge, implant placement becomes more predictable. Three-dimensional radiographic assessment can also improve the accuracy of guided implant surgery and help avoid surgical complications. The use of digital planning technologies has further increased the precision of implant positioning and prosthetic rehabilitation [4]. Another important component of comprehensive examination is the evaluation of occlusion. Excessive occlusal forces, bruxism, and unfavorable bite relationships can lead to mechanical overload of implants and prosthetic components. Therefore, functional analysis should be carried out before definitive implant treatment. In selected patients, occlusal correction, protective night guards, or changes in prosthetic design may be necessary to decrease excessive loading.



Laboratory tests may also be required in patients with systemic diseases or suspected metabolic abnormalities. Blood glucose levels, coagulation parameters, vitamin D status, and other indicators may influence surgical safety and healing. Such investigations are not mandatory for every patient, but they are important when the medical history suggests an increased risk of complications.

Comprehensive examination also improves communication between the clinician and the patient. The patient should receive clear information about the expected treatment stages, possible complications, alternative options, healing period, costs, and the importance of oral hygiene. Informed consent and realistic expectations are important elements of successful implant therapy. Long-term implant survival depends not only on successful osseointegration but also on appropriate maintenance after treatment. Regular professional examinations, radiographic control, plaque management, and monitoring of peri-implant tissues are necessary to detect early signs of inflammation or bone loss. Evidence shows that supportive periodontal and peri-implant care can significantly reduce the risk of biological complications [5].

In summary, comprehensive patient examination before dental implantation provides the basis for safe, predictable, and individualized treatment. Medical history assessment, clinical examination, radiological diagnostics, evaluation of bone and soft tissues, occlusal analysis, and identification of systemic and local risk factors are all essential components of implant planning. Careful assessment reduces the likelihood of complications and increases the probability of long-term functional and aesthetic success.

References:

[1] Buser D., Sennerby L., De Bruyn H. Modern implant dentistry based on osseointegration: 50 years of progress, current trends and open questions. *Periodontology* 2000. 2017;73(1):7–21.

[2] Chrcanovic B.R., Albrektsson T., Wennerberg A. Smoking and dental implants: A systematic review and meta-analysis. *Journal of Dentistry*. 2015;43(5):487–498.

[3] Derks J., Tomasi C. Peri-implant health and disease: A systematic review of current epidemiology. *Journal of Clinical Periodontology*. 2015;42(Suppl. 16)–S171.

[4] Bornstein M.M., Scarfe W.C., Vaughn V.M., Jacobs R. Cone beam computed tomography in implant dentistry: A systematic review focusing on guidelines, indications, and radiation dose risks. *International Journal of Oral & Maxillofacial Implants*. 2014;29(Suppl.):55–77.

[5] Schwarz F., Derks J., Monje A., Wang H.L. Peri-implantitis. *Journal of Clinical Periodontology*. 2018;45(Suppl. 20)–S266.