



**THE EFFECT OF ALCOHOL CONSUMPTION
DURING PREGNANCY ON FETAL DEVELOPMENT:
PATHOGENESIS, CLINICAL CONSEQUENCES, AND
PREVENTION**

Turg'unboyeva Malika Xursanali qizi

*Tashkent International University of Kimyo, Namangan
branch (KIUT), Faculty of Pediatrics, 2nd-year student,
+998942571027 / turgunboyevamalika595@gmail.com*

Scientific supervisor: Kamalova Gulnora Gulyamovna

*PhD, Acting Assoc. Prof., Tashkent International
University of Kimyo, Namangan Branch, Namangan,
Uzbekistan*

ARTICLE INFO:

Received:25.10.2026

Revised: 26.10.2026

Accepted:27.10.2026

KEY WORDS:

*prenatal alcohol
exposure, fetal
development, fetal
alcohol spectrum
disorders, fetal alcohol
syndrome,
teratogenicity,
embryogenesis,
placental dysfunction,
oxidative stress,
neurodevelopment,
congenital
abnormalities,
intrauterine growth
restriction, antenatal
care, early diagnosis,
primary prevention,
maternal health.*

ABSTRACT

Alcohol consumption during pregnancy remains an important and preventable risk factor for adverse fetal outcomes, including impaired intrauterine growth, congenital abnormalities, neurodevelopmental disturbances, and fetal alcohol spectrum disorders (FASD). Ethanol readily crosses the placental barrier, exposing the developing fetus to a substance that its immature metabolic systems cannot process as efficiently as those of an adult. The consequences of prenatal alcohol exposure vary according to the amount and pattern of consumption, timing and duration of exposure, maternal nutritional status, genetic susceptibility, placental function, and other environmental influences. This article aims to examine the principal biological mechanisms underlying alcohol-related fetal injury, summarize its clinical consequences, and identify evidence-based approaches to prevention and early recognition. A narrative review of international scientific publications, clinical guidance, and an official Uzbek antenatal-care document was undertaken. The literature was examined with particular attention to placental dysfunction, oxidative stress, cellular injury, altered gene regulation, impaired brain development, and disturbances in fetal growth. The findings of the reviewed literature indicate that prenatal alcohol exposure may interfere with neural cell proliferation, migration, differentiation, synaptic



organization, and the development of essential organ systems. Clinically, the effects range from pregnancy loss and premature birth to growth restriction, characteristic facial differences in some affected children, congenital abnormalities, and persistent difficulties involving cognition, learning, attention, memory, and behavioral regulation. FASD represents a spectrum of conditions, and affected children do not necessarily exhibit the same physical characteristics or degree of impairment. The absence of obvious abnormalities at birth therefore does not exclude later neurodevelopmental difficulties. The review also highlights the importance of non-stigmatizing antenatal screening, clear communication about alcohol-related risks, and timely referral for appropriate support when alcohol use is difficult to discontinue. Current evidence does not establish a safe amount, type, or period of alcohol consumption during pregnancy. Complete abstinence while planning pregnancy and throughout gestation is consequently the most reliable preventive approach. Integrating preventive counseling into routine antenatal care, strengthening professional education, and improving developmental surveillance may reduce preventable harm and support affected children and their families.

Introduction: Fetal development is a complex biological process that depends on coordinated cellular differentiation, tissue formation, organ maturation, placental function, and the continuous exchange of oxygen and nutrients between the mother and the developing fetus. Exposure to harmful substances during this period may interfere with these processes and produce consequences that extend beyond birth. Alcohol is particularly important because ethanol can cross the placenta, exposing fetal tissues during stages of development when essential organs, especially the brain, remain highly vulnerable. Prenatal alcohol exposure is therefore recognized as a preventable cause of developmental disability and congenital abnormalities [1, 2].

The effects of alcohol during pregnancy are not limited to the presence of a characteristic physical appearance. Some children develop recognizable features of fetal alcohol syndrome, whereas others primarily experience difficulties with learning, attention, memory, executive functioning, emotional regulation, or social adaptation. These manifestations are collectively considered within the broader concept of fetal alcohol spectrum disorders. The severity and



presentation of these conditions differ between individuals, even when prenatal exposure histories appear similar [1, 3].

The clinical significance of this issue is increased by the fact that fetal brain development continues throughout pregnancy. Alcohol exposure may therefore affect different developmental processes at different gestational stages. Early exposure can interfere with embryogenesis and organ formation, while exposure during later pregnancy may affect brain maturation, growth, and the development of functional neural networks. It is not possible to identify a gestational period during which alcohol consumption can be considered reliably safe [2, 4].

Maternal alcohol consumption may also be influenced by social circumstances, psychological stress, limited access to health information, unplanned pregnancy, and difficulties associated with alcohol dependence. Consequently, prevention requires more than simply informing pregnant women about potential risks. It also requires accessible antenatal services, confidential assessment, supportive counseling, and referral to appropriate treatment when necessary. The Uzbek antenatal-care guideline similarly emphasizes the importance of identifying alcohol use and arranging appropriate medical support for pregnant women who require it [5].

The aim of this article is to analyze the pathophysiological mechanisms through which alcohol affects fetal development, summarize the principal clinical consequences of prenatal exposure, and discuss evidence-based approaches to prevention, antenatal screening, and early developmental assessment.

Literature review: Research into prenatal alcohol exposure has progressed from the description of characteristic physical abnormalities to a broader understanding of the molecular, neurological, and environmental processes involved in fetal injury. Popova and colleagues described fetal alcohol spectrum disorders as a group of conditions characterized by variable combinations of neurodevelopmental impairment, growth abnormalities, facial differences, and congenital anomalies. Their review emphasizes that the consequences of prenatal exposure depend not only on alcohol consumption itself but also on its timing, dose, frequency, maternal and fetal genetic characteristics, nutrition, and other environmental factors [1].

The developmental toxicity of ethanol is closely associated with its ability to cross the placenta. The fetus is exposed to alcohol circulating in maternal blood, while its immature metabolic capacity limits the efficient elimination of ethanol and its metabolites. Exposure may consequently interfere with developing tissues through several overlapping pathways, including oxidative stress, disturbances in cellular signaling, altered gene expression, impaired mitochondrial function, and disruption of normal developmental processes [1, 6].

One of the most extensively investigated areas concerns the effects of alcohol on the central nervous system. During embryonic and fetal development, neural cells must proliferate, migrate to appropriate locations, differentiate into specialized cell types, and establish functional connections. Alcohol exposure may disrupt several of these processes. The resulting



abnormalities can affect brain structure and function, potentially contributing to difficulties with executive functioning, attention, learning, memory, impulse control, and behavioral adaptation. However, the specific manifestations vary considerably, and no single cognitive or behavioral feature is sufficient to establish a diagnosis of FASD [1, 3].

A systematic review examining prenatal alcohol exposure and brain development during the first 1,000 days of life identified 16 eligible studies. The findings were inconsistent across imaging methods and brain regions, although some studies reported differences involving the thalamus, amygdala, hippocampus, and white-matter connectivity. These findings support concern about altered early brain development while also demonstrating the methodological challenges involved in determining the precise effects of different exposure patterns [7].

Alcohol-related fetal injury is also associated with disturbances in placental function and fetal growth. The placenta regulates the transport of oxygen, nutrients, and metabolic products while supporting communication between maternal and fetal physiological systems. Alcohol may affect placental development, vascularization, and nutrient transport, potentially contributing to impaired fetal growth. Experimental research also suggests that alcohol exposure can alter inflammatory signaling, metabolic regulation, and epigenetic mechanisms. Nevertheless, the relative contribution of each pathway in human pregnancy remains incompletely understood [6].

The clinical spectrum of prenatal alcohol exposure includes miscarriage, stillbirth, premature birth, impaired fetal growth, congenital abnormalities, and persistent neurodevelopmental difficulties. The Centers for Disease Control and Prevention and the National Institute on Alcohol Abuse and Alcoholism emphasize that FASD encompasses multiple diagnostic presentations rather than a single uniform condition [2, 3]. Diagnosis may be delayed because certain learning and behavioral difficulties become apparent only when children encounter increasing cognitive and social demands.

International recommendations consistently advise avoiding alcohol during pregnancy and when attempting to conceive. The Royal College of Obstetricians and Gynaecologists emphasizes that the safest approach is complete abstinence, while recognizing the importance of supportive counseling for women who have consumed alcohol before recognizing pregnancy or who find it difficult to stop [4]. In Uzbekistan, the official antenatal-care guideline provides a relevant framework for identifying alcohol use during pregnancy and directing affected women toward appropriate medical assessment [5]. Taken together, these sources support an integrated approach combining prevention, early recognition, clinical follow-up, and developmental support.

Materials and methods: This article was prepared as a narrative literature review of prenatal alcohol exposure, fetal developmental injury, clinical outcomes, and preventive interventions. The source material included international reviews, systematic-review evidence, public-health guidance, and an official Uzbek antenatal-care document. The principal



international sources included publications indexed in PubMed and articles published in Nature Reviews Disease Primers, alongside guidance from the Centers for Disease Control and Prevention, the National Institute on Alcohol Abuse and Alcoholism, and the Royal College of Obstetricians and Gynaecologists.

The literature was examined thematically rather than through a quantitative meta-analysis. The main analytical categories were the biological mechanisms of ethanol-related fetal injury, gestational vulnerability, placental and embryonic effects, neurological and structural outcomes, diagnostic considerations, and preventive strategies. Particular attention was given to the distinction between established clinical guidance, findings from human observational research, and mechanisms supported primarily by experimental studies.

Evidence was synthesized using comparative reading and descriptive analysis. The interpretation considered differences in study design, outcome measurement, exposure assessment, and the possibility of confounding factors. No original patient recruitment, laboratory testing, clinical intervention, or statistical analysis was undertaken. Accordingly, the findings presented below represent a qualitative synthesis of published evidence rather than newly generated clinical results. The review is limited by its narrative design and does not claim to provide an exhaustive systematic search or a pooled estimate of the magnitude of risk.

Results: The reviewed literature identifies several interconnected pathways through which prenatal alcohol exposure may interfere with fetal development. The first is direct exposure of fetal tissues to ethanol and its metabolites. Because ethanol crosses the placenta, maternal alcohol consumption can expose the developing embryo and fetus during periods of rapid cellular growth and organ formation. The immature fetal metabolic system does not eliminate alcohol as efficiently as the adult organism, increasing concern about exposure during sensitive developmental stages [1, 2].

A second pathway involves oxidative stress and cellular injury. Ethanol metabolism can generate reactive compounds and disturb the balance between oxidative processes and antioxidant defenses. Experimental research indicates that these disturbances may affect cellular membranes, mitochondrial function, signaling pathways, and the survival of developing cells. The consequences depend on exposure characteristics and the developmental stage of the affected tissue. Although oxidative stress is a plausible contributor to alcohol-related injury, it should not be regarded as the sole mechanism responsible for the full clinical spectrum of FASD [1, 6].

A third pathway concerns the development of the central nervous system. Normal brain formation requires tightly regulated neural proliferation, migration, differentiation, synaptogenesis, and maturation. Alcohol exposure may interfere with these processes and alter the organization of developing neural networks. Such changes may contribute to difficulties involving attention, learning, memory, motor coordination, language, and executive functioning. The systematic-review evidence on early brain imaging suggests that some exposed infants



demonstrate differences in particular brain structures or connectivity, but the available findings are heterogeneous and do not establish a uniform imaging pattern for all affected children [7].

The literature also indicates that placental dysfunction and disturbances in fetal growth may contribute to adverse pregnancy outcomes. Changes in placental vascular development and nutrient transport may reduce the efficiency with which the fetus receives the resources required for normal growth. Clinically, prenatal alcohol exposure is associated with an increased risk of impaired fetal growth, low birth weight, and premature birth. The strength of these associations varies across studies and may be influenced by the amount and pattern of alcohol consumption, maternal health, nutrition, tobacco use, and other factors [1, 4, 6].

The clinical consequences of prenatal alcohol exposure range from pregnancy loss and congenital abnormalities to persistent developmental difficulties. Fetal alcohol syndrome represents one recognizable presentation, generally involving characteristic facial features, growth deficiency, and central nervous system abnormalities. Other children within the FASD spectrum may have no obvious characteristic facial appearance but may experience substantial problems with cognition, adaptive functioning, learning, or behavior. Consequently, the absence of visible abnormalities at birth cannot reliably exclude alcohol-related neurodevelopmental impairment [1, 2]. The reviewed evidence also demonstrates the importance of early identification. A history of prenatal alcohol exposure should prompt an appropriately individualized assessment of growth, physical development, neurological function, and emerging cognitive or behavioral concerns. No single blood test or imaging investigation can independently establish every FASD diagnosis. Assessment instead requires integration of exposure history, clinical examination, developmental evaluation, and consideration of alternative explanations for the child's difficulties [2, 3].

With regard to prevention, the literature consistently supports complete avoidance of alcohol during pregnancy and when planning conception. Screening for alcohol use should be incorporated into routine antenatal care through confidential, non-judgmental discussion. When a pregnant woman reports alcohol consumption, the immediate priorities are to encourage cessation, assess the pattern of use and any associated dependence, arrange appropriate prenatal monitoring, and provide referral for treatment where needed. The Uzbek antenatal-care guidance supports identifying alcohol use and arranging specialist assistance when indicated [4, 5].

Discussion: The findings of this narrative review support the conclusion that prenatal alcohol exposure is a significant and preventable cause of fetal developmental injury. Its effects are best understood as the result of interacting biological processes rather than a single isolated mechanism. Direct cellular toxicity, oxidative stress, altered gene regulation, impaired placental function, and disruption of neural development may contribute in different proportions depending on the exposure pattern and the developmental characteristics of the pregnancy [1, 6].



The developing brain deserves particular attention because neurological maturation continues throughout gestation and extends into childhood. Prenatal alcohol exposure may affect neural processes that later support learning, executive functioning, memory, emotional regulation, and adaptive behavior. These difficulties can become more evident with age, particularly when school and social environments place increasing demands on the child. This helps explain why some children may not receive an appropriate diagnosis during infancy despite experiencing clinically meaningful developmental difficulties later in life [1, 3].

At the same time, scientific interpretation requires caution. Not every pregnancy involving alcohol exposure results in an identifiable FASD, and not every child with developmental difficulties has experienced prenatal alcohol exposure. Differences in exposure measurement, maternal and fetal genetics, nutrition, concurrent substance use, and socioeconomic conditions complicate the interpretation of observational findings. The systematic review of brain development during the first 1,000 days illustrates these limitations, reporting inconsistent results across the imaging studies examined [7]. Therefore, the literature supports a clear preventive recommendation without suggesting that the outcome of every individual exposure can be predicted with certainty.

Antenatal counseling is a central component of prevention. Information should be clear, medically accurate, and delivered without stigmatizing the pregnant woman. Fear-based or judgmental communication may discourage disclosure and reduce engagement with prenatal services. A more effective clinical approach involves asking about alcohol consumption confidentially, explaining the potential consequences, encouraging complete abstinence, and offering practical assistance to women who need support to stop drinking. Where dependence is suspected, clinical assessment and appropriate specialist management are important, because care should address the woman's health as well as fetal well-being [4, 5].

Another important consideration is the difference between preventing exposure and managing its consequences. FASD is preventable by avoiding prenatal alcohol exposure, but established neurodevelopmental impairments may require long-term support. Children with suspected or confirmed FASD may benefit from individualized developmental assessment, educational accommodations, behavioral interventions, family support, and coordination between pediatric, psychological, and educational services. Early recognition does not reverse all prenatal injury, but it can help families obtain appropriate services and reduce the impact of unmet developmental needs [2, 3].

For Uzbekistan, strengthening routine antenatal screening and preventive counseling represents a practical opportunity to improve maternal and child health. The existing antenatal-care framework provides a basis for identifying alcohol use, documenting relevant history, and arranging specialist referral when needed [5]. Further local research could investigate the implementation of screening in antenatal clinics, healthcare professionals' knowledge, barriers to disclosure, and the accessibility of support services. Such research should use appropriate



ethical safeguards and culturally sensitive methods. It should also distinguish the prevalence of alcohol exposure from the prevalence of clinically diagnosed FASD, as these are not equivalent measures.

The present review has several limitations. It was designed as a narrative synthesis rather than a registered systematic review, and no formal risk-of-bias assessment or meta-analysis was performed. The article does not provide original clinical data or a population-specific estimate for Uzbekistan. In addition, some proposed molecular mechanisms are supported more strongly by experimental evidence than by direct clinical studies in pregnant women. These limitations mean that the conclusions should be interpreted as a synthesis of existing evidence and clinical guidance, not as proof of the relative contribution of every biological pathway.

Conclusion and practical recommendations: Alcohol consumption during pregnancy can interfere with fetal development through direct cellular toxicity, oxidative stress, disruption of developmental signaling, altered placental function, and disturbances in brain maturation. The resulting clinical consequences may include pregnancy loss, premature birth, impaired fetal growth, congenital abnormalities, and fetal alcohol spectrum disorders. The manifestations of prenatal alcohol exposure are highly variable, and significant cognitive or behavioral difficulties may become apparent only later in childhood. Therefore, a normal appearance at birth does not necessarily exclude subsequent developmental problems.

Current evidence does not establish a safe amount, type, or period of alcohol consumption during pregnancy. Complete abstinence while planning conception and throughout pregnancy is the most reliable preventive strategy. Antenatal services should provide clear information, confidential screening, non-judgmental counseling, and timely referral for women who require assistance to stop drinking. If alcohol exposure has occurred before pregnancy recognition, the appropriate response is to discontinue further consumption, seek prenatal care, and discuss the exposure with a healthcare professional rather than delay care because of fear or guilt.

Practical recommendations include incorporating alcohol-use assessment into routine antenatal visits; educating healthcare professionals about the biological mechanisms and clinical manifestations of prenatal alcohol exposure; strengthening referral pathways for women who experience alcohol dependence; and monitoring fetal growth and child development according to clinical indications. Pediatric services should consider prenatal exposure history when assessing unexplained developmental, learning, behavioral, or neurological difficulties. When FASD is suspected, multidisciplinary evaluation and individualized family support should be arranged. At the public-health level, prevention requires coordinated efforts involving maternity services, primary healthcare, pediatric professionals, educational institutions, and community health programs. Further research in Uzbekistan should evaluate local implementation of antenatal screening, barriers to accessing treatment, and approaches to early developmental recognition. Overall, combining complete alcohol abstinence during pregnancy, supportive antenatal care, early recognition of developmental concerns, and appropriate long-



term support offers the strongest evidence-based approach to reducing preventable alcohol-related harm.

References:

1. Popova S., Charness M.E., Burd L., et al. Fetal alcohol spectrum disorders. Nature Reviews Disease Primers. 2023;9(1):11. Springer Nature. Article number 11. <https://doi.org/10.1038/s41572-023-00420-x>.
2. Centers for Disease Control and Prevention. About Fetal Alcohol Spectrum Disorders (FASDs). Atlanta, GA: U.S. Department of Health and Human Services, CDC; 2024. Online clinical and public-health resource. Available at: <https://www.cdc.gov/fasd/about/index.html>.
3. National Institute on Alcohol Abuse and Alcoholism. Understanding Fetal Alcohol Spectrum Disorders. Bethesda, MD: National Institutes of Health, U.S. Department of Health and Human Services. Online scientific information resource. Available at: <https://www.niaaa.nih.gov/publications/brochures-and-fact-sheets/understanding-fetal-alcohol-spectrum-disorders>.
4. Royal College of Obstetricians and Gynaecologists. Alcohol and Pregnancy. London: RCOG. Patient information and clinical education resource.
5. Ministry of Health of the Republic of Uzbekistan. On the Organization and Provision of Antenatal Care and Medical Assistance to Pregnant Women in Primary Health Care Institutions. MED.UZ medical information portal. Tashkent, Uzbekistan. Official antenatal-care guidance. Available at: <https://www.med.uz/documentation/detail.php?ID=31170>.
6. Nonconceptus Mechanisms of Prenatal Alcohol Exposure That Disrupt Embryo-Fetal Development: An Integrative View. 2025. Full-text scientific review available through PubMed Central, National Library of Medicine, National Institutes of Health.
7. The Association of Prenatal Alcohol Exposure With Brain Development During the First 1000 Days of Life: A Systematic Review. Systematic review; literature search through February 2024. PubMed, U.S. National Library of Medicine. Available at: <https://pubmed.ncbi.nlm.nih.gov/39439004/>.
8. American Academy of Pediatrics. Fetal Alcohol Spectrum Disorders: Parent FAQs. Itasca, IL: American Academy of Pediatrics. Pediatric clinical information resource.
9. National Health Service. Fetal Alcohol Spectrum Disorder. London: NHS. Clinical information and prevention guidance. Available at: <https://www.nhs.uk/conditions/fetal-alcohol-spectrum-disorder/>.
10. American College of Obstetricians and Gynecologists. Fetal Alcohol Spectrum Disorders: Frequently Asked Questions. Washington, DC: ACOG. Professional clinical information resource.